

Splitting and focusing of neutrino collective states

Mattias Marklund^{†‡}, Padma K. Shukla^{§||} and Lennart Stenflo^{||}

[†] Department of Electromagnetics, Chalmers University of Technology, SE-412 96 Göteborg, Sweden

[§] Institut für Theoretische Physik IV, Fakultät für Physik und Astronomie, Ruhr-Universität Bochum, D-44780 Bochum, Germany

^{||} Department of Plasma Physics, Umeå University, SE-901 87 Umeå, Sweden

Abstract. It is shown that the collective nonlinear interactions between intense neutrino or anti-neutrino fluxes and a dense neutrino plasma are governed by a multi-dimensional coupled cubic Schrödinger equation in which the interaction potential is positive or negative depending on the neutrino type. The cubic Schrödinger equation describes the splitting and focusing of intense neutrino beams due to the nonlinear excitations associated with the modifications of the individual neutrino energies in a dense neutrino background.

Keywords: neutrino properties, supernova neutrinos, ultra high energy photons and neutrinos

Neutrinos and anti-neutrinos are produced in the Sun and in supernovae during violent processes. The propagation of intense neutrinos in ionised media (dense plasmas) is a topic of current interest in view of the fact that neutrinos may be massive [1]. Because of their weak interaction with charged and neutral particles, neutrinos (or anti-neutrinos) can travel great distances without being affected appreciably by material obstacles. Massive neutrinos can account for neutrino oscillations and associated flavour change in solar plasmas [2, 3], while in supernovae and cosmology they may play a decisive role in reviving shocks for supernovae explosions, as well as in accounting for the missing dark matter in our Universe [4].

In the present paper, we employ a semi-classical approach to consider the nonlinear interaction between neutrinos and anti-neutrinos in neutrino plasmas. Using an effective field theory approach, a system of coupled nonlinear Schrödinger equations is derived, and it is shown that these equations admit splitting and focusing of intense neutrino (or anti-neutrino) beams due to the nonlinearities associated with the weak nuclear forces of the neutrinos/anti-neutrinos that are interacting with the background neutrinos.

[‡] E-mail: marklund@elmagn.chalmers.se

Suppose a single neutrino (or anti-neutrino) moves in a neutrino–anti-neutrino admixture. The energy E of the neutrino (anti-neutrino) is then given by (see, e.g. [6, 7, 8])

$$E = \sqrt{p^2 c^2 + m^2 c^4} + V_{\pm}(\mathbf{r}, t), \quad (1)$$

where $\mathbf{p}$ is the neutrino (anti-neutrino) momentum, c is the speed of light in vacuum, and m is the neutrino mass. The effective potential for a neutrino moving on a background of its own flavor is [7, 10] (see also [9, 11]) $V_{\pm}(\mathbf{r}, t) = \pm 2\sqrt{2}G_F(n - \bar{n})$, while the potential for a neutrino moving on a background of a different flavour takes the form $V_{\pm}(\mathbf{r}, t) = \pm\sqrt{2}G_F(n - \bar{n})$. Here $G_F/(\hbar c)^3 \approx 1.2 \times 10^{-5} \text{ GeV}^{-2}$, where G_F is the Fermi constant, n ($\bar{n}$) is the density of the background neutrinos (anti-neutrinos), and the $+$ ($-$) is for the propagating neutrino (anti-neutrino). From Eq. (1) we can, using standard techniques [12], obtain a Schrödinger equation for a neutrino (anti-neutrino) wave packet $\Psi(\mathbf{r}, t)$ [13] (see also Ref. [14] for a similar treatment of neutrino–electron interactions)

$$i\frac{\partial\Psi}{\partial t} + \frac{\hbar}{2m_{\text{eff}}}\nabla^2\Psi - \frac{V_{\pm}}{\hbar}\Psi = 0, \quad (2)$$

where $m_{\text{eff}} = m\gamma$, $\gamma = (1 - v^2/c^2)^{-1/2}$, and v is the magnitude of the group velocity of the neutrino wave packet.

In the case of self-interacting neutrinos and anti-neutrinos, moving with the same group velocity, we have $n = \langle |\Psi_+|^2 \rangle$ and $\bar{n} = \langle |\Psi_-|^2 \rangle$, where Ψ_+ and Ψ_- are the neutrino and anti-neutrino wave functions, respectively, and the angular bracket denotes the ensemble average. Thus, in this case Eq. (2) turns into the coupled system

$$i\frac{\partial\Psi_{\pm}}{\partial t} + \frac{\alpha}{2}\nabla^2\Psi_{\pm} \mp \beta(\langle |\Psi_+|^2 \rangle - \langle |\Psi_-|^2 \rangle)\Psi_{\pm} = 0, \quad (3)$$

where $\alpha = \hbar/m_{\text{eff}}$ and $\beta = 2\sqrt{2}G_F/\hbar$ for neutrinos of the same flavor.

Suppose now that the neutrino pulses are coherent and uni-directional. If one species, the pump species (denoted by the index p), dominates the number density as compared to the signal species (denoted by the index s), i.e. $n_p \gg n_s$, we obtain the equation

$$i\frac{\partial\Psi_s}{\partial t} + \frac{\alpha}{2}\frac{\partial^2\Psi_s}{\partial z^2} + \beta|\Psi_p|^2\Psi_s = 0, \quad (4a)$$

for the signal wave function Ψ_s , where the driving nonlinear term is determined from

$$i\frac{\partial\Psi_p}{\partial t} + \frac{\alpha}{2}\frac{\partial^2\Psi_p}{\partial z^2} - \beta|\Psi_p|^2\Psi_p = 0. \quad (4b)$$

It is well known that Eq. (4b) have modulationally stable dark soliton solutions [15], of which the fundamental soliton is

$$\Psi_p = \Psi_0 \tanh\left(\sqrt{\beta/\alpha}\Psi_0 z\right) \exp(-i\beta\Psi_0^2 t). \quad (5)$$




Figure 1. Contour plot of $\ln[n_s(t, z)/n_{s0}]$ ($b = 1$) for the splitting case.




Figure 2. Contour plot of $\ln[n_s(t, z)/n_{s0}]$ ($b = 1$) for the focusing case.

Following Ref. [16], we perform a perturbation analysis of Eq. (4a) for a given solution $\Psi_p(t, z)$ to Eq. (4b). Separating $\Psi_s(t, z)$ into its real amplitude $\sqrt{n_s(t, x)}$ and phase $\theta_s(t, z)$, we have, to lowest order in t ,

$$\theta_s(t, z) \approx \beta \int_0^t |\Psi_p(t', z)|^2 dt' \approx \beta |\Psi_p(0, z)|^2 t, \quad (6)$$

where the last equality follows if the interaction takes place over a time-scale short compared to the time over which the pump amplitude changes. The signal number

density $n_s(t, z)$ is then

$$n_s(t, z) \approx n_s(0, z) \exp \left[-\frac{\alpha \beta t^2}{2} \left(\frac{d^2 n_p(0, z)}{dz^2} + \frac{dn_p(0, z)}{dz} \frac{d}{dz} \ln n_s(0, z) \right) \right], \quad (7)$$

where $n_p(t, z) = |\Psi_p(t, z)|^2$ is the pump number density. Using the fundamental soliton (5) as the pump pulse, with $\Psi_0^2 = n_p(0, 0) \equiv n_{p0}$, and a Gaussian distribution for the initial signal pulse,

$$n_s(0, z) = n_{s0} \exp(-z^2/a^2), \quad (8)$$

where a is the width and $n_{s0} = n_s(0, 0)$, we find that

$$n_s(t, z) = n_{s0} \exp \left\{ -b^2 \zeta^2 - \left[(3 \operatorname{sech}^2 \zeta - 2) \operatorname{sech}^2 \zeta - 2b^2 \tanh \zeta \operatorname{sech}^2 \zeta \right] \tau^2 \right\}, \quad (9)$$

where $\zeta = z/z_0$, $\tau = \alpha t/z_0^2$, $b = z_0/a$, and $z_0 = (\beta n_{p0}/\alpha)^{1/2}$. We note that b , τ and ζ are dimensionless. The solution (9) is plotted in Fig. 1, where the splitting of the collective neutrino state can clearly be seen.

In the case of neutrino–neutrino or anti-neutrino–anti-neutrino interaction, Eq. (3) will be modified according to

$$i \frac{\partial \Psi_i}{\partial t} + \frac{\alpha}{2} \nabla^2 \Psi_i - \beta (\langle |\Psi_1|^2 \rangle + \langle |\Psi_2|^2 \rangle) \Psi_i = 0, \quad (10)$$

where $i = 1, 2$. For a dark soliton pump beam (5) and a Gaussian signal beam (8), the approximate dynamics, analogous to the neutrino–anti-neutrino interaction, can be obtained from (9) by letting $\tau^2 \rightarrow -\tau^2$. The result is seen in Fig. 2, where the focusing is depicted.

The above discussion is based on the approximate solution (9). However, numerical studies of the system (4a) and (4b) show the same behaviour, when (5) and (8) are used as initial conditions. Thus, a more complete solution supports the notion that the approximate analytical expression (9) gives an accurate description of the dynamics of interacting neutrinos and anti-neutrinos.

In conclusion, we have considered the nonlinear interactions between neutrinos and anti-neutrinos, mediated by a dense neutrino background of the same flavour. It is shown that the self-interaction between neutrinos and anti-neutrinos comes about due to their coupling via the weak nuclear force. Such interactions are governed by coupled cubic Schrödinger equations for the slowly varying envelopes of the neutrino and anti-neutrino wave functions. Analytical and numerical analysis of the multi-dimensional cubic Schrödinger equations reveal that a collective neutrino state can be split or focused due to the presence of a dense soliton background. The present results suggest that nonlinear excitations in neutrino plasmas can significantly affect the propagation of intense neutrino and anti-neutrino beams. The neutrino beam splitting can account for the energy loss of neutrinos, while the self-focusing renders neutrino burst propagation over a long distance.

References

- [1] Ahmad Q R, *et al.* 2002 *Phys. Rev. Lett.* **89** 011301
- [2] Bahcall J 1989 *Neutrino Astrophysics* (Cambridge: Cambridge University Press)
- [3] Raffelt G 1996 *Stars as Laboratories for Fundamental Physics* (Chicago: Chicago University Press)
- [4] Dolgov A D 2002 *Phys. Rep.* **370** 333
- [5] Sofue Y and Rubin V C 2001 *Ann. Rev. Astron. Astrophys.* **39** 137
- [6] Bethe H A 1987 *Phys. Rev. Lett.* **56** 1305; Bethe H A 1990 *Rev. Mod. Phys.* **62**, 801
- [7] Kuo T K and Pantaleone J 1989 *Rev. Mod. Phys.* **61** 937
- [8] Silva L O, *et al.* 2000 *Astrophys. J.* **127** 481
- [9] Weldon H A 1982 *Phys. Rev. D* **26** 2789
- [10] Nötzhold D and Raffelt G 1988 *Nucl. Phys. B* **307** 924
- [11] Nunokawa H, *et al.* 1997 *Nucl. Phys. B* **501** 17
- [12] Hasegawa A 1975 *Plasma Instabilities and Nonlinear Effects* (Berlin: Springer-Verlag)
- [13] Marklund M, *et al.* 2003 submitted[astro-ph/0306013].
- [14] Tsintsadze N L, *et al.* 1998 *Phys. Plasmas* **5** 3512; Shukla P K, *et al.* 2002 *Phys. Plasmas* **9** 3625
- [15] Kivshar Y S and Luther-Davies B 1998 *Phys. Rep.* **298** 81
- [16] Helczynski L, *et al.* 2000 *Physica Scripta* **T84** 81